

# Solutions to Sheet 12

## Exercise 1

Let  $A$  be a ring and let  $G$  be a finite group acting on  $A$  by ring automorphisms. Let  $A^G$  be the ring of invariants of  $G$  in  $A$ .

1. Show that  $A$  is integral over  $A^G$ .
2. Assume that  $A$  is a domain with quotient field  $K$ . Show that  $K^G = \text{Quot}(A^G)$ .

### Solution.

1. Let  $x \in A$  be any element. Then

$$P_x(T) = \prod_{g \in G} (T - g(x))$$

is a monic polynomial with coefficients invariant under  $G$  (by symmetry). As  $P_x(x) = 0$ ,  $x$  is integral over  $A^G$ , and we are done.

2.  $G$  acts on  $K$  via  $g(\frac{x}{y}) = \frac{g.x}{g.y}$ . One readily verifies  $\text{Quot}(A^G) \subseteq K^G$ . For the other inclusion, assume  $\frac{x}{y} \in K^G$ . Now we can write

$$\frac{x}{y} = \frac{x \prod_{\text{id} \neq h \in G} h(y)}{\prod_{h \in G} h(y)}.$$

The denominator is  $G$ -invariant. But now the numerator is as well, as

$$x \prod_{\text{id} \neq h \in G} h(y) = \frac{x}{y} \prod_{h \in G} h(y)$$

is a product of two  $G$ -invariant elements. Hence  $\frac{x}{y}$  can be expressed as the quotient of two  $G$ -invariant objects. I struggled way too much with this exercise.

## Exercise 2

1. Let  $A$  be a normal domain with quotient field  $K$  and let  $G$  be a finite group acting on  $A$  by ring automorphisms. Show that  $A^G$  is normal.
2. Let  $k$  be a field of characteristic  $\neq 2$ . Show that  $k[x, y, z]/(z^2 - xy)$  is normal.

### Solution.

1. This is just collecting what we did in exercise 1.  $A$  is algebraically closed in its quotient field  $K$ . Also,  $A$  is integral over  $A^G$  and  $K$  is integral over  $K^G = \text{Quot}(A^G)$ . But now  $K$  is integral over  $A^G$ , in particular  $K^G \subseteq K$  is integral over  $A^G$ .
2. We have  $k[x, y, z]/(z^2 - xy) \cong k[x^2, xy, y^2] = k[x, y]^G$ , where  $G = \{\pm 1\}$  acts via

$$(-1).f(x, y) = f(-x, -y).$$

Now we are in the situation of part 1, and as  $k[x, y]$  is normal, we are done.

### Exercise 3

Let  $L/K$  be a finite Galois extension of number fields with Galois group  $G$ . Show that  $\mathcal{O}_L$  is stable under action of  $G$  and that  $\mathcal{O}_L^G = \mathcal{O}_K$ .

**Solution.** To show that  $G$  is invariant under the action of  $G$ , let  $x \in \mathcal{O}_L$  be an element with  $f(x) = 0$ , where  $f \in \mathcal{O}_K[T]$  is monic and irreducible. Let  $\sigma \in \text{Gal}(L/K)$ . Write  $f^\sigma$  for the polynomial that arises when applying sigma to the coefficients of  $f$ . Now  $f^\sigma(\sigma x) = \sigma(f(x)) = 0$ . (I just realized we have  $f^\sigma = f$ . Whatever.)

We now show the second statement. By Galois theory, we know that  $L^G = K$ . As  $\mathcal{O}_L \subseteq L$  this shows  $\mathcal{O}_L^G = \mathcal{O}_L \cap L^G$ . By definition,  $\mathcal{O}_L$  is the integral closure of  $\mathcal{O}_K$  in  $\mathcal{O}_L$ . This directly shows  $\mathcal{O}_L^G \supseteq \mathcal{O}_K$ . The other direction follows because every element in  $\mathcal{O}_L \cap K$  is integral over  $\mathcal{O}_K$ , which (by the definition of the integral closure) implies that  $\mathcal{O}_L \cap K \subseteq \mathcal{O}_K$ .

### Exercise 4

Let  $k$  be a field and let  $A := k[x, y]/(y^2 - x^3 - x^2)$ .

1. Show that  $A$  is a domain.
2. Show that  $t = y/x \in \text{Quot}(A)$  does not lie in  $A$ .
3. Show that  $t$  is integral over  $A$ .
4. Show that  $\text{Quot}(A) = k(t)$  and that  $k[t] \subseteq \text{Quot}(A)$  is the normalization of  $A$ .

### Solution.

1. We have

$$k(x)[y]/(y^2 - x^2(x+1)) \cong k(x)[y]/((y/x)^2 - (x+1)),$$

and this is a quadratic field extension. In particular,  $(y^2 - x^3 - x^2)$  is irreducible in  $k(x)[y]$ , hence also irreducible in  $k[x, y]$ . Alternatively, the Eisenstein criterion over  $k[x]$  works.

2. Suppose  $x/y \in A$ . Now we have  $(x, y) = (x)$ . But in  $k[x, y]/(y^2 - x^3 - x^2)$ , we have  $y \notin (x)$ .
3. We have  $t^2 = \frac{y^2}{x^2} = x + 1 \in A$ .
4. What does this even mean?! The normalization is simply the integral closure of  $A$  in  $\text{Quot}(A)$ . First, note  $k(t) \subseteq \text{Quot}(A)$  because  $t \in \text{Quot}(A)$ . For the reverse statement, note that the calculation in part 1 shows that

$$\text{Quot}(A) \subseteq k(x)[y]/(y^2 - x^3 - x^2) = k(x)[t]/(t^2 - x - 1) = k(t).$$

Let  $N \subseteq \text{Quot}(A)$  denote the normalization of  $A$ . We have  $N \supseteq k[t]$  because  $t$  is integral over  $A$ , and  $N \subseteq k[t]$  because  $k[t]$  is integrally closed in  $k(t) = \text{Quot}(A)$ .